

2022 Nishina Memorial Prize

Dr. Eiji Saitoh

Professor, Graduate School of Engineering, University of Tokyo



“Pioneering contribution to the physics of spin current”

The charge of an electron and its flow, the electric current, have always been the main physical quantities of interest in electronics. The angular momentum, or the spin, is another fundamental physical quantity of an electron that generates magnetic moment through its polarization and plays an important role in the physics of magnetism and related engineering. Spintronics has emerged as a research field that seeks novel physical phenomena and functionalities through the control of spins, where the flow of spins, or the spin current, has attracted particular interest.

The spins of conduction electrons in a conductor usually point either upward or downward in a 50-50 ratio, and thus the spin transfer associated with the flow of electrons is averaged out to zero. The balance between the up- and down-spin flows can be changed to yield a net spin current, e.g., by the flow of spin-polarized electrons in magnetic materials and by the "spin Hall effect", in which, due to spin-orbit interaction in the material, electrons are subject to a force perpendicular to the current direction depending on their spin orientations. However, a direct method for measuring the spin current with an external probe was elusive, and experiments were limited to indirect estimation, e.g., through the observation of "spin accumulation" generated at the sample edge by the spin current.

Dr. Saitoh discovered the "inverse spin Hall effect" in 2006 as a scheme for the direct measurement of spin current [1]. He used a bilayer metal film of platinum (Pt) and ferromagnetic alloy, Permalloy ($\text{Ni}_{81}\text{Fe}_{19}$). When spin excitations were generated in the Permalloy layer through ferromagnetic resonance and injected into the Pt probe layer through the interface, the spin current was converted into electric current due to the strong spin-orbit interaction in Pt, and a voltage signal was detected between the two ends of the Pt layer along the direction perpendicular to the magnetic field. This enabled direct measurement of the spin current for the first time and led to the vast development of related research.

Since then, Dr. Saitoh and his colleagues have discovered various physical phenomena involving spin currents by using this spin-current detection technique mentioned above. A major achievement among them is the "spin Seebeck

effect", in which a spin current is generated by a temperature gradient applied to a magnetic material and is injected into a probe electrode, e.g., made from Pt, to produce a voltage through the inverse spin Hall effect [2]. In conventional thermoelectric devices based on the Seebeck effect, two different conductors are combined in parallel under a thermal gradient, and the voltage is generated from the difference in the density of states and scattering properties of the conduction carriers in each conductor. The clever idea that led to the spin Seebeck effect was to utilize the difference in the behavior of the two spin states of electrons in a single magnetic material.


Dr. Saitoh and his collaborators then extended the physics of spin current from conductors to insulators. They showed that the spin current, as a flow of angular momentum in solids, is carried not only by conduction electrons but also by spin excitations in insulators, greatly expanding the concept of spintronics. In ferromagnetic insulating oxides such as yttrium iron garnet (YIG), spin waves, which are collective excitation modes of ordered spins of localized electrons, or magnons as their elementary excitations, propagate over long distances without being scattered by conduction electrons. In their 2010 paper, Dr. Saitoh *et al.* reported that they injected a spin current into a YIG thin film by using the spin Hall effect in a Pt electrode and observed the propagation of the spin current using the inverse spin Hall effect in another Pt electrode [3]. This result also revealed that angular momentum is transferred between conduction electrons in the metal electrodes and spin excitations in the insulator through the exchange interaction at the interface. Dr. Saitoh *et al.* also observed the spin Seebeck effect in YIG-related oxides [4], pioneering research on thermoelectric devices using insulating materials. Furthermore, they experimentally demonstrated that spin currents are carried not only by magnons in ferromagnets but also by various elementary excitations in solids, such as magnons in antiferromagnets, spinons in quantum spin liquids [5], magnons in nuclear spin wave modes [6], and magnetic polarons, coupled excitations of magnons and phonons. These achievements revealed the broad potential of spin current as a probe for the study of the physical properties of materials.

References:

- [1] E. Saitoh, M. Ueda, H. Miyajima, and G. Tatara, "Conversion of spin current into charge current at room temperature: inverse spin-Hall effect", *Applied Physics Letters* **88**, 182509 (2006).
- [2] K. Uchida, S. Takahashi, K. Harii, J. Ieda, W. Koshibae, K. Ando, S. Maekawa, and E. Saitoh, "Observation of the spin Seebeck effect", *Nature* **455**, 778 (2008).
- [3] Y. Kajiwara, K. Harii, S. Takahashi, J. Ohe, K. Uchida, M. Mizuguchi, H. Umezawa, H. Kawai, K. Ando, K. Takanashi, S. Maekawa, and E. Saitoh, "Transmission of electrical signals by spin-wave interconversion in a magnetic insulator", *Nature* **464**, 262 (2010).
- [4] K. Uchida, J. Xiao, H. Adachi, J. Ohe, S. Takahashi, J. Ieda, T. Ota, Y. Kajiwara, H. Umezawa, H. Kawai, G. E. W. Bauer, S. Maekawa, and E. Saitoh, "Spin Seebeck insulator", *Nature Materials* **9**, 894 (2010).
- [5] D. Hirobe, M. Sato, T. Kawamata, Y. Shiomi, K. Uchida, R. Iguchi, Y. Koike, S. Maekawa, and E. Saitoh, "One-dimensional spinon spin currents", *Nature Physics* **13**, 30 (2017).
- [6] Y. Shiomi, J. Lustikova, S. Watanabe, D. Hirobe, S. Takahashi, and E. Saitoh, "Spin pumping from nuclear spin waves", *Nature Physics* **15**, 22 (2019).

Dr. Eiichiro Komatsu

Director, Max Planck Institute for Astrophysics



“Contribution to the standard cosmology based on cosmic microwave background”

As a theory that explains the global homogeneity and isotropy of the universe, inflationary cosmology, which postulates that the universe experienced exponential accelerated expansion long before primordial nucleosynthesis, is a very attractive idea. However, in order to establish it as the standard cosmology, it is important not only to explain such qualitative observational facts but also to quantitatively verify its predictions. These include that the universe is spatially flat, has an almost scale-invariant spectrum originating from quantum fluctuations, and generates adiabatic curvature fluctuations and tensor fluctuations (quantum gravitational waves) that approximately follow Gaussian statistics.

Dr. Komatsu, under the advice of Professor Spergel at Princeton University, focused on the statistical nature of fluctuations and developed a methodology to quantitatively evaluate the deviation from the Gaussian distribution using cosmic microwave background radiation. While Gaussian distributions can be characterized only by their amplitude and variance, non-Gaussian distributions have infinite possibilities, making it difficult to quantitatively evaluate deviations from a Gaussian distribution. They proposed to constrain the deviation to a single parameter, the nonlinear parameter, and to limit it by observations of the cosmic microwave background radiation [1]. Dr. Komatsu established a methodology to measure this by using three-point correlations (bispectrum) of the cosmic microwave background radiation and applied it to the data observed by the COBE (COsmic Background Explorer) [2].

He also joined the research team of the Wilkinson Microwave Anisotropy Probe (WMAP), which was in progress at the time, and applied it to the first-year observation data, quantitatively verifying for the first time in the world that the nonlinear parameter has no significant finite value and the curvature fluctuation is consistent with a Gaussian distribution. This was the first quantitative verification in the world [3]. At the same time, they also performed a detailed analysis of the spectrum of the fluctuations and found the existence of nearly scale-invariant adiabatic curvature fluctuations, as predicted by standard inflationary cosmology.

WMAP also showed that the spatial curvature of the universe is below the limit of measurement, verifying that

space is flat as predicted by inflationary cosmology. WMAP also measured the amount of cold dark matter and dark energy, which is responsible for accelerating the expansion of the universe, including error estimates, and found that the current universe consists of about 5% baryons, 22% cold dark matter, and 73% dark energy, although these values were updated by the subsequent observation of the Planck mission. In addition, the Hubble parameter, which expresses the expansion rate of the universe, was precisely measured, and it was revealed for the first time that the universe is 13.7 billion years old. These results confirmed the validity of the cold dark matter model with a cosmological term, which was obtained at the end of the 20th century based on a variety of observations. It can be said that cosmology was promoted to the precision science for the first time with the achievement of WMAP. In the third year of the WMAP, Dr. Komatsu took charge of polarization analysis in addition to his responsibilities in the first year, and in the fifth year, he became responsible for the entire analysis.

During this period, the system of each cosmological parameter measured gradually improved, and significant values were measured for the spectral index, which represents the deviation of the curvature fluctuation from the scale invariance of the spectrum. Theoretically, the standard single-field slow-roll inflation model predicts that the spectral exponent is slightly less than 1, and its value is a major clue to identifying the model. Signs of a spectral index deviating from 1 began to appear in the fifth year of WMAP data, but were rejected with a 99.5% statistical confidence level in the seventh year of data [4].

As described above, WMAP played a decisive role in establishing today's standard cosmology, which predicts that structure formation occurred in a universe filled with cold dark matter and dark energy, with curvature fluctuations that are nearly scale-invariant and follow a Gaussian distribution as the initial conditions as predicted by inflationary cosmology. Dr. Komatsu played a leading role in the analysis of WMAP data and contributed greatly to the contemporary standard cosmology.

References

- [1]. E. Komatsu and D. N. Spergel “Acoustic signatures in the primary microwave background bispectrum”, *Physical Review D* **63**, 063002 (2001).
- [2]. E. Komatsu, B. D. Wandelt, D. N. Spergel, A. J. Banday and K. M. Górski, “Measurement of the cosmic microwave background bispectrum on the COBE DMR sky maps”, *Astrophysical Journal*, **566**, 19 (2002).
- [3]. E. Komatsu, et al. “First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Tests of Gaussianity”, *Astrophysical Journal Supplement Series*, **148**, 119 (2003).
- [4]. E. Komatsu, et al. “Seven Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological interpretation”, *Astrophysical Journal Supplement Series*, **192**, 18 (2011).